




REMEDY GUIDE · HERBS

# Turmeric buying without the colour trap

Bright powder tells you almost nothing about strength, safety or dose. This guide follows turmeric from monsoon fields through drying and extraction so you can read assays and origin papers with calm confidence.

- **Root powder strength**  
Whole ground rhizome usually tests near 2 to 9 percent curcuminoids.
- **Concentrated extract power**  
Refined extracts often assay near 80 to 95 percent curcuminoids.
- **Papers before colour**  
Lot assays plus lead screening tell more than any golden shade.

By Kathryn Foster · Updated Sep 23, 2026 · 8 min read

CONTENTS

|    |                                                |    |                                                 |
|----|------------------------------------------------|----|-------------------------------------------------|
| 01 | Field work to dry rhizome                      | 05 | Reasons makers pair extract with pepper and fat |
| 02 | Drying choices that alter hue and curcumin     | 06 | Contaminant checks plus provenance files        |
| 03 | Contents of milled whole root                  | 07 | The checklist                                   |
| 04 | Concentration into strong curcuminoid extracts | 08 | Sources and further reading                     |

---

A single lot assay can change what you buy more than shade, aroma or story. Ground rhizome naturally carries only a small share of curcuminoids, while refined extracts carry a far larger share, and only paperwork shows which jar you hold.

01

Field work to dry rhizome

Turmeric comes from *Curcuma longa* L., a perennial herb that forms underground rhizomes and belongs to Zingiberaceae. Growers raise it as a seasonal crop across warm humid zones, with large plantings in India and China. India continues as the leading grower, so many traceability sheets lead back to its districts.

Lifting takes place at maturity, after leafy shoots wither and underground weight reaches its peak. Farm crews dig finger and mother pieces, rinse soil away, then boil, cut or polish them in line with regional custom. Cutting has practical value because slender slices lose water quicker and more uniformly than intact fingers.

Newly lifted rhizomes hold abundant moisture near 70 to 80 percent. One nutrient summary lists water around 80 to 90 percent, then carbohydrates near 13 percent, proteins near 2 percent, minerals near 2 percent and lipids below 1 percent. Such water weight makes fresh rhizomes bulky and quick to spoil.

Drying lowers water activity to restrain enzymes, microbes and spoilage chemistry and to cut shipping weight. Commerce therefore moves dried stock, whether intact dried fingers, milled powder or feedstock for concentration. A parcel that remains moist can develop mould and uneven character before encapsulation ever begins.

---

## 02

### Drying choices that alter hue and curcumin

Drying does more than drive off water. Warmth, illumination, airflow and duration shift hue, phenolic levels, curcumin levels and volatile oil recovery. A six way trial examined sun drying, convection oven drying, vacuum oven drying, infrared drying, microwave drying and freeze drying using 3 mm slices of turmeric.

Under those trial conditions freeze drying kept the clearest yellow tone, the smallest browning score at 17.07, phenolics at 78.90 mg GAE per g, curcumin at 3.83 percent and volatile oil at 4 percent. Cool handling without light kept both pigment and aromatic oil intact. High capital cost leaves this route uncommon for bulk spice volumes.

Sun drying remains the most common practice. It brings long exposure, variable batches and chance of microbial growth. The sharpest fall in curcuminoids was linked to ultraviolet breakdown. Extended time in open yards fades colour while dampness departs slowly.

Other heat based routes fall in the middle. Convection, vacuum, infrared and microwave systems shorten cycles and curb microbial hazard relative to open yards. Each supplies heat differently, so pigment keeping and oil yield change with temperature, pressure and cut thickness. Retail packs seldom name dryer type, though bulk vendors should retain it.

#### FROM THE EDITORS

Ask for the lab assay, not only the colour lore A vivid yellow may come from gentle drying, from added pigment or only from cultivar traits, while a duller powder may still test inside its normal curcuminoid interval.

## 03

### Contents of milled whole root

Milled root retains the full rhizome matrix. That matrix holds carbohydrates, protein, minerals, fibre, volatile and fixed oils, alongside yellow pigments known as curcuminoids. Whether that broader matrix with turmerones and oils acts differently from separated curcuminoids remains unproven.

Curcuminoids can reach as much as 10 percent of dried turmeric mass. The mixture centres on curcumin, demethoxycurcumin and bisdemethoxycurcumin, reported between 2 and 9 percent in turmeric. Another account places whole root powder near 2 to 8 percent curcuminoid by weight, while a culinary account cites near 3 to 8 percent curcumin with season.

The sums stay modest. A typical 3 g dessert spoon of powder supplies roughly 30 to 90 mg curcumin on average. Kitchen custom generally adds about 0.5 to 3 g powder daily in meals. Culinary intake therefore provides tens of milligrams rather than the hundreds of milligrams assessed in extract research.

Bench assays can measure curcumin, demethoxycurcumin and bisdemethoxycurcumin separately. One survey across six Indian sites refluxed 15 g portions in a Soxhlet with 150 mL ethylene dichloride at 70 degrees C for 6 hours, then read total curcuminoids by absorbance close to 420 nm and single curcuminoids by HPLC. Absorbance close to 420 nm reports a total, whereas HPLC resolves the trio. Precise harvest timing and district values for any retail jar call for lot level assay documents.

---

## 04

### Concentration into strong curcuminoid extracts

Curcumin repels water and will not dissolve in it, though it dissolves in dimethyl sulfoxide, acetone, ethanol and oil. Plain water extracts little pigment from powder. Producers rely on food grade solvents, warmth and contact time to enrich the markers.

Classic solvent recovery of curcumin with 70 percent hydroethanolic liquid attained 69.1 percent recovery. Longstanding isolation routes include alcoholic Soxhlet work, hexane treatment followed by acetone work, and hot ethanol filtration with kerosene and petroleum ether stages that produce orange recrystallized curcumin needles. Those belong in laboratories and factories rather than home kitchens.

Recovery responds to liquid choice, warmth, flow and dwell. Ethanol assisted subcritical water work at 90 degrees C, 4 mL per min and 25 wt percent ethanol returned 4.12 wt percent dry basis yield, above Soxhlet at 2.71 wt percent and ultrasound assisted plus ordinary solvent work near 0.85 wt percent. Modest shifts in settings move the amount of curcuminoid drawn from the matrix.

Typical trade extracts are normalised to roughly 80 to 95 percent total curcuminoids. Whole root at 2 to 8 percent and refined extract toward 95 percent stand far apart. A 500 mg capsule at 95 percent carries near 475 mg curcuminoids, while 500 mg of 3 percent powder carries near 15 mg. Improved uptake systems further cloud dose math because equal milligrams may behave unlike one another across technologies.

---

## 05

### Reasons makers pair extract with pepper and fat

Concentrated curcuminoid extracts present a further hurdle beyond strength. Uptake is limited and clearance is swift, so brands include piperine from black pepper or combine curcuminoids with oils and phospholipids. The panel reveals which strategy a maker selected.

In human research, 20 mg piperine taken with 2,000 mg curcumin raised measured bioavailability near 2,000 percent. Piperine blocks UDP glucuronosyltransferase enzymes and P glycoprotein, retarding curcumin breakdown and limiting return of curcumin from gut cells into the lumen. Slower conjugation across gut wall and liver keeps more parent material circulating for a period.

Fat based formats follow separate reasoning. Since curcumin favours oil, firms suspend extract in oils, micelles, liposomes or phospholipid pairings. The aim is to place a water shy powder in a lipid friendly carrier. Study labels, patents and brand terms differ, so milligram to milligram matching is hard without human results on that precise format.

Serving ranges mirror the divide between cooking and capsules. Knee osteoarthritis research often tests near 500 to 1,500 mg daily of curcuminoid extract. Home cooking remains near 0.5 to 3 g powder daily in dishes. Greater potency does not automatically mean greater safety because liver injury reports cluster around products engineered for stronger uptake. Extended safety of potent enhanced uptake formulas past a few months of study remains unestablished.

#### FROM THE EDITORS

Request a certificate of analysis for the lot. Home water and palm checks described in blogs are not laboratory tests for lead or curcuminoid level, so treat them as curiosity rather than proof.

## 06

### Contaminant checks plus provenance files

Milled spice brings hazards that refining can partly sidestep and partly retain. Earth, irrigation water, open drying grounds and mill gear may introduce heavy metals. Pigment fraud is a separate intentional issue.

Lead chromate, a poisonous yellow colourant, is added by certain vendors to deepen turmeric powder hue, especially across bulk chains. A 2014 Harvard University survey detected lead up to 483 ppm in turmeric from 18 rural Bangladesh households, against a permitted level of 2.5

ppm. Stanford investigators found no sign of tainted turmeric outside Bangladesh and observed that tighter import scrutiny pushed many large Bangladesh processors to end or curb lead chromate practice.

Pesticide traces are a worry in milled spice, yet no quantities are given for the home jar. That silence matters to homes shopping for a partner, child or parent. Provenance files can shrink doubt even where they cannot erase it.

Purity checks named for shoppers include USDA Organic, NSF or USP review, and outside review by ConsumerLab or Labdoor, alongside counsel to obtain batch level heavy metal sheets. Organic addresses cultivation methods, while NSF, USP, ConsumerLab and Labdoor address testing schemes with varied scope. None substitutes for a lot level certificate of analysis.

Ask the maker for the bottle lot code, the nation and district of growth, the drying technique where recorded, the analytical method and outcome for total curcuminoids, the HPLC split of the trio for extracts, and the heavy metal table with thresholds. A founder able to name growers and display sheets earns confidence. Missing files do not establish a poor product, yet they keep the chain unclear.

---

## 07

### The checklist

- 
- ☐ Fit format to purpose: culinary root powder or measured curcuminoid extract
- 
- ☐ Judge total curcuminoids by percent rather than hue or narrative
- 
- ☐ Seek the trio on record: curcumin, demethoxycurcumin, bisdemethoxycurcumin
- 
- ☐ Calculate milligrams per capsule and per day from label percent
- 
- ☐ Record piperine level if shown, commonly around 20 mg in research
- 
- ☐ Record oil or phospholipid carrier and resist cross brand milligram math
- 
- ☐ Review Supplement Facts and companion ingredients for carriers and pepper
- 
- ☐ Require lot matched certificate of analysis naming assay technique
- 
- ☐ Obtain batch level heavy metal sheet with lead before buying.
- 
- ☐ Seek USDA Organic alongside NSF, USP, ConsumerLab or Labdoor review
- 
- ☐ File nation, harvest lot and drying route where disclosed
- 
- ☐ Treat potent uptake formulas with caution and raise them with a clinician
- 

## 08

### Sources and further reading

- Optimizing drying techniques for turmeric (*Curcuma longa* L.): impacts on color, curcumin, and essential oil composition · [pmc.ncbi.nlm.nih.gov](https://pubmed.ncbi.nlm.nih.gov/)
- Impacts of turmeric and its principal bioactive curcumin on human health: Pharmaceutical, medicinal, and food applications: A comprehensive · [pmc.ncbi.nlm.nih.gov](https://pubmed.ncbi.nlm.nih.gov/)
- Optimization of ethanol-modified subcritical water extraction of curcuminoids from turmeric (*Curcuma longa* L.) rhizomes: Comparison with con · [sciencedirect.com](https://www.sciencedirect.com/)
- Turmeric Whole Root vs Standardized Curcumin for Everyday Wellness and Targeted Joint Support · [suplmnt.app](https://suplmnt.app)
- Evaluation of Curcuminoids in Turmeric Rhizome (*Curcuma Longa* L.) Collected from Different Places in India · [biotech-asia.org](https://biotech-asia.org)
- Different Methods of Isolation of Curcumin · [biosmartnotes.com](https://biosmartnotes.com)

For information only, not medical advice. Herbs and supplements can interact with medicines and are not right for everyone, especially in pregnancy; check with a doctor or pharmacist before you start one.

---



WRITTEN BY

**Kathryn Foster**

Kathryn runs the news desk and the remedy guides at Root to Bottle. She keeps a shelf of old herbals at her desk and checks every traditional use against what the trials actually found, then writes down both.

---